

Hip 63510C, Hip 73786B, and nine new isolated high proper motion T dwarf candidates from UKIDSS DR6 and SDSS DR7

R.-D. Scholz

Astrophysikalisches Institut Potsdam, An der Sternwarte 16, 14482 Potsdam, Germany
e-mail: rdscholz@aip.de

Received 16 February, 2010; accepted 18 March, 2010

ABSTRACT

Aims. Completing the poorly known substellar census of the solar neighbourhood, especially with respect to the coolest brown dwarfs, will lead to a better understanding of failed star formation processes and binary statistics with different environmental conditions.

Methods. Using UKIDSS data and their cross-correlation with the SDSS, we searched for high proper motion mid- to late-T dwarf candidates with extremely blue near-infrared ($J-K < 0$) and very red optical-to-near-infrared ($z-J > +2.5$) colours.

Results. With 11 newly found T dwarf candidates, the proper motions of which range between 100 and 800 mas/yr, we increased the number of UKIDSS T dwarf discoveries by $\approx 30\%$. Large proper motions were also measured for six of eight previously known T4.5-T9 dwarfs detected in our survey. All new candidates can be classified as T5-T9 dwarfs based on their colours. Two of these objects were found to be common proper motion companions of Hipparcos stars with accurate parallaxes. The latter allow us to determine absolute magnitudes from which we classify Hip 63510C as T7 and Hip 73786B as T6.5 dwarfs with an uncertainty of ± 1 spectral subtype. The projected physical separation from their low-mass (M0.5 and K5) primaries is in both cases about 1200 AU. One of the Hipparcos stars has already a known very low-mass star or brown dwarf companion on a close astrometric orbit (Hip 63510B = Gl 494B). With distances of only 11.7 and 18.6 pc, deduced from their primaries respectively for Hip 63510C and Hip 73786B, various follow-up observations can easily be carried out to study these cool brown dwarfs in more detail and to compare their properties with those of the already well-investigated primaries.

Key words. Astrometry – Proper motions – Stars: distances – Stars: kinematics and dynamics – brown dwarfs – solar neighbourhood

1. Introduction

New deep near-infrared surveys like UKIDSS¹ and the Canada-France Brown Dwarf Survey (CFBDS; Delorme et al. 2008b) provide a powerful tool for detecting even more of the coolest brown dwarfs (mid- and late-T dwarfs) in the solar neighbourhood than previously found in the SDSS (Abazajian et al. 2009) and 2MASS (Skrutskie et al. 2006). From 155 currently known T dwarfs (see Gelino et al. 2009 and references therein), there were 52 discovered with SDSS and 49 with 2MASS during the last 12 years. The number of discoveries from the UKIDSS large area survey (LAS) has already grown to 33 in only three years, where the majority were found by Lodieu et al. (2007) and Pinfield et al. (2008). The latest-type objects discovered in the SDSS and 2MASS are of spectral type T7 (Chiu et al. 2006) and T8 (Burgasser et al. 2002; Tinney et al. 2005; Looper, Kirkpatrick, & Burgasser 2007) respectively, whereas a few T8.5-T9 dwarfs have been found in UKIDSS (Warren et al. 2007; Burningham et al. 2008, 2009) and CFBDS (Delorme et al. 2008a).

Since most of the stellar neighbours of the Sun were originally detected as high proper motion stars, one can expect to find the nearest cool brown dwarfs in new high proper motion surveys using deep multi-epoch near-infrared imaging data over large areas of sky, as e.g. eventually provided by two epochs of

J-band data in UKIDSS. However, one can already try to use the SDSS (Abazajian et al. 2009), the deepest available optical survey overlapping with the ongoing UKIDSS LAS, as the first epoch. The nearest cool brown dwarfs may still be visible in the red optical z-band of the SDSS data observed a few years before the UKIDSS observations started.

We describe a high proper motion survey for nearby T dwarfs using mainly UKIDSS and SDSS data which led to the discovery of 11 new mid- to late-T dwarf candidates, including two common proper motion objects of nearby red dwarf stars with accurate Hipparcos parallaxes.

2. Selection of T dwarf candidates from UKIDSS

For the selection of T dwarf candidates, we used the available cross-matching with the SDSS DR7 (Abazajian et al. 2009) as provided in the UKIDSS data base. To select faint objects that appear very red in optical-to-near-infrared colours, but blue in the near-infrared, as expected for mid- to late-T dwarfs, we used the following criteria:

$$J1 > 11, \quad J1 - K < 0, \quad z - J1 > +2.5,$$

where $J1$ are the first epoch J-band data, and z comes from the nearest matching SDSS object within the search radius of 10 arcsec. By doing so, we allowed for missing Y- and H-band data. Whereas our main search criterion was the negative $J-K$ colour, we excluded all kind of optically blue objects (early-type stars and white dwarfs) with the $z-J$ colour cut. Late-M and L dwarfs would also meet the $z-J$ colour but not the $J-K$ colour

¹ The UKIDSS project is defined in Lawrence et al. (2007). UKIDSS uses the UKIRT Wide Field Camera (WFCAM; Casali et al. 2007) and a photometric system described in Hewett et al. (2006) which is situated in the Mauna Kea Observatories (MKO) system (Tokunaga et al. 2002). The pipeline processing and science archive are described in Irwin et al. (2010) and Hambly et al. (2008).

criterion. L subdwarfs (e.g. Lodieu et al. 2010) have negative $J-K$ but bluer $z-J$ ($\approx +1.5$). However, low-metallicity T dwarfs would possibly also satisfy our colour cuts, as the peculiar T6 dwarf 2MASS J0937+29 (Burgasser et al. 2006a) has $z-J \approx +3.5$ (derived from SDSS DR7 and Leggett et al. 2010 photometry). Note that a large $i-z$ was not required as a pre-condition. 4940 targets were found satisfying these criteria.

UKIDSS $Y, J1, J2, H$, and K finding charts of all these candidates were inspected to exclude the vast majority of them consisting of ghost images of bright stars, images affected by diffraction spikes, satellite trails, galaxies, and asteroids. For the remaining star-like sources, the corresponding SDSS finding charts and DR7 data were checked for possible counterparts. We have also used the SDSS database available at Princeton (Finkbeiner et al. 2004), where we found some additional epochs for determining the proper motions. Only 11 new candidates with a unique very red SDSS counterpart (detected only in the z -band with $z \approx 20$, whereas $i > 23$ values listed in the SDSS DR7 indicate that there was no flux above the noise) within a smaller search radius of 5 arcsec and a significant proper motion were finally selected and are presented in Sect. 3. Note that these counterparts cannot be background late-M or L dwarfs, since they would show up in UKIDSS with $J \approx 18$ and $K \approx 16-17$, which is not the case. In addition, eight previously known T4.5-T9 dwarfs were detected and proper motions determined for six of them with identified SDSS counterparts (Sect. 4).

We have also checked 2MASS, DENIS (Epchtein et al. 1997) and SSS (Hambly et al. 2001) I -band data for possible counterparts but failed to find any for the new T dwarf candidates due to their faintness.

3. Astrometry & photometry of T dwarf candidates

Full names and photometry of 11 new T dwarf candidates are listed in Table 1. All candidates have $ugriz$ data where only the z magnitudes are at or below the SDSS detection limit, whereas the $ugri$ measurements are clearly above these limits. According to Abazajian et al. (2009), a 95% detection repeatability for point sources is provided at $u=22.0$, $g=22.2$, $r=22.2$, $i=21.3$, $z=20.5$.

For the proper motion determination we took advantage of the sometimes slightly different epochs between the $YJHK$ -band measurements (up to one year epoch differences) and of the additional epochs (separated by several years from the UKIDSS epochs) provided by the SDSS (more details can be found in Tables 2 and 4). For the proper motion solutions we used linear fitting over all available epochs combined with a search for




Fig. 1. 3×3 arcmin² UKIDSS J -band images, left: ULAS J1300+12 (= Hip 63510C) with the bright primary, Hip 63510A in the upper left corner, right: ULAS J1504+05 (= Hip 73786B) with its primary, Hip 73786A on the right side. North is up, East to the left.




Fig. 2. 2×2 arcmin² UKIDSS J -band images of ULAS J0329+04, ULAS J0819+21, ULAS J0945+07 (top row, from left to right), ULAS J1012+10, ULAS J1417+13, ULAS J1449+11 (middle row), ULAS J2320+14, ULAS J2321+13, and ULAS J2342+08 (bottom row). North is up, East to the left.

the correct counterparts at the expected positions in data from additional epochs.

The errors of the proper motion components given in Tables 3, 5, and 6 are estimated from the scatter around the best-fit line in each coordinate, and describe the quality of the fit. Due to the small number of available epochs, these formal errors may be unrealistically small (few data points may be well-aligned by chance) compared to the true errors. If we know the errors in the positions of the objects measured at different epochs in the SDSS and UKIDSS, we can estimate which error in their proper motions would be expected for the given epoch differences using error propagation.

The SDSS DR7 statistical errors per coordinate for bright stars are 45 mas, with systematic errors of less than 20 mas (Abazajian et al. 2009). However, for faint objects at the SDSS survey limit, the astrometric accuracy is limited by photon statistics to about 100 mas (Pier et al. 2003). The UKIDSS astrometric accuracy varies between 50 mas at low galactic latitudes and 100 mas at high galactic latitudes (Lawrence et al. 2007). For our objects, mostly located at intermediate galactic latitudes ($|b|=40^\circ-50^\circ$), we assume uncertainties of 100 mas and 70 mas respectively for single SDSS and UKIDSS positions, which include smaller colour-dependent systematic errors in the astrometry expected to be of the order of a few tens of milliarcsec. We combine all available SDSS data of a given object in a mean first epoch and all UKIDSS data in a mean second epoch, where the accuracy of the mean positions is getting higher with the square root of the number of available single epochs, and compute the expected proper motion errors from error propagation. These errors (ϵ) are given in the last column of Tables 3, 5, and 6. For the previously known T dwarfs (Table 6), available 2MASS and DENIS positions were also assigned 100 mas individual errors and used together with the SDSS positions for the mean first epoch, which was then compared with the mean UKIDSS position to compute ϵ .

Table 4. Multi-epoch positions α, δ (J2000.0) of isolated objects

object			epoch	ID	source
<i>s</i>	<i>'</i>	<i>''</i>	yr		
ULAS J0329+04:					
20.1535	30	24.390	2005.781	5714	SDSS DR7
20.1809	30	24.415	2006.816	6485	SDSS DR7
20.2211	30	24.558	2008.784	2336494	UKIDSS <i>H</i>
20.2227	30	24.538	2008.784	2336510	UKIDSS <i>K</i>
20.2170	30	24.549	2008.805	2338708	UKIDSS <i>Y</i>
20.2207	30	24.583	2008.805	2338724	UKIDSS <i>J</i>
ULAS J0819+21:					
18.5794	03	11.246	2004.209	4508	SDSS DR7
18.5888	03	10.809	2007.049	1184392	UKIDSS <i>J</i>
18.5848	03	10.480	2008.907	2340123	UKIDSS <i>H</i>
18.5792	03	10.497	2008.907	2340143	UKIDSS <i>K</i>
18.5849	03	10.441	2008.910	2341325	UKIDSS <i>Y</i>
18.5809	03	10.430	2008.910	2341345	UKIDSS <i>J</i>
ULAS J0945+07:					
16.1702	55	46.304	2002.194	3031	SDSS DR7
16.3889	55	45.586	2007.052	1186762	UKIDSS <i>J</i>
16.3974	55	45.632	2007.052	1188655	UKIDSS <i>J</i>
16.4008	55	45.652	2007.060	1343254	UKIDSS <i>H</i>
16.3905	55	45.650	2007.060	1343620	UKIDSS <i>H</i>
16.3977	55	45.636	2007.063	1330616	UKIDSS <i>K</i>
16.3937	55	45.589	2007.063	1332824	UKIDSS <i>K</i>
16.3947	55	45.584	2007.066	1335120	UKIDSS <i>Y</i>
16.3919	55	45.569	2007.066	1336848	UKIDSS <i>Y</i>
ULAS J1012+10:					
43.6322	21	04.108	2002.953	3538	SDSS DR7
43.5471	21	01.826	2007.005	1147818	UKIDSS <i>K</i>
43.5350	21	01.683	2007.249	1354063	UKIDSS <i>H</i>
43.5350	21	01.698	2007.304	1415926	UKIDSS <i>J</i>
43.5356	21	01.698	2007.304	1417555	UKIDSS <i>Y</i>
ULAS J1417+13:					
56.2478	30	45.565	2003.472	3996	SDSS DR7
56.2275	30	45.875	2007.367	1450483	UKIDSS <i>Y</i>
56.2263	30	45.868	2007.367	1450579	UKIDSS <i>J</i>
56.2226	30	45.943	2008.222	2318417	UKIDSS <i>K</i>
56.2236	30	45.915	2008.222	2318429	UKIDSS <i>H</i>
ULAS J1449+11:					
01.9706	47	12.307	2003.409	3971	SDSS DR7
01.9050	47	11.390	2007.260	1367513	UKIDSS <i>J</i>
01.9064	47	11.415	2007.260	1368654	UKIDSS <i>Y</i>
01.9138	47	11.267	2007.293	1402270	UKIDSS <i>K</i>
01.9024	47	11.389	2007.293	1402702	UKIDSS <i>H</i>
ULAS J2320+14:					
35.1020	48	28.996	2000.735	1739	SDSS P
35.1064	48	28.978	2000.910	1904	SDSS P
35.2839	48	29.835	2007.729	1716958	UKIDSS <i>K</i>
35.2900	48	29.815	2007.729	1716859	UKIDSS <i>H</i>
35.2914	48	29.815	2007.729	1717061	UKIDSS <i>Y</i>
35.2890	48	29.836	2007.729	1717217	UKIDSS <i>J</i>
ULAS J2321+13:					
23.7591	54	59.058	2000.735	1739	SDSS P
23.7724	54	58.799	2000.910	1904	SDSS P
23.7910	54	58.261	2001.639	2507	SDSS P
23.7990	54	54.895	2007.666	1689556	UKIDSS <i>J</i>
23.7990	54	54.943	2007.666	1690721	UKIDSS <i>K</i>
23.7990	54	54.918	2007.666	1690865	UKIDSS <i>Y</i>
23.7969	54	54.935	2007.666	1691223	UKIDSS <i>H</i>
ULAS J2342+08:					
28.9358	56	20.123	2006.715	6354	SDSS DR7
28.9455	56	20.133	2007.901	1830486	UKIDSS <i>H</i>
28.9423	56	20.140	2007.901	1830585	UKIDSS <i>K</i>
28.9659	56	20.111	2008.732	2335155	UKIDSS <i>Y</i>
28.9655	56	20.105	2008.732	2335171	UKIDSS <i>J</i>

Note: The ID is the run or multiframe number for SDSS or UKIDSS respectively. SDSS P stands for the SDSS data base in Princeton.

Table 5. Proper motions of isolated objects in mas/yr

Object	$\mu_{\alpha} \cos \delta$	μ_{δ}	ε
ULAS J0329+04	+323 $\pm$ 15	+60 $\pm$ 07	32
ULAS J0819+21	+5 $\pm$ 14	-168 $\pm$ 07	24
ULAS J0945+07	+685 $\pm$ 13	-142 $\pm$ 07	21
ULAS J1012+10	-327 $\pm$ 11	-558 $\pm$ 06	25
ULAS J1417+13	-76 $\pm$ 03	+77 $\pm$ 03	24
ULAS J1449+11	-242 $\pm$ 21	-244 $\pm$ 18	27
ULAS J2320+14	+387 $\pm$ 05	+121 $\pm$ 02	11
ULAS J2321+13	+56 $\pm$ 15	-577 $\pm$ 10	10
ULAS J2342+08	+229 $\pm$ 55	-9 $\pm$ 09	48

Notes: Proper motions come from linear fitting of the positions given in Table 4. ε are the expected errors (see text).

Table 6. Proper motions of known T dwarfs in mas/yr

Object	SpT (Ref)	$\mu_{\alpha} \cos \delta$	μ_{δ}	ε
SDSS J0325+04	T5.5 (1)	-185 $\pm$ 06	+15 $\pm$ 07	27
2MASS J0755+22	T5.0 (2)	-13 $\pm$ 03	-241 $\pm$ 05	11
SDSS J0830+01	T4.5 (2)	+214 $\pm$ 06	-329 $\pm$ 09	08
2MASS J1231+08	T5.5 (2)	-1185 $\pm$ 09	-1050 $\pm$ 12	14
SDSS J1504+10	T7.0 (1)	+338 $\pm$ 13	-350 $\pm$ 11	21
2MASS J2339+13	T5.0 (2)	+380 $\pm$ 05	-971 $\pm$ 05	12

Notes: Proper motions as obtained from linear fitting of UKIDSS, SDSS, and (if available) 2MASS and DENIS positions. ε are the expected errors (see text). For full names and references, see Gelino et al. (2009). Spectral type (SpT) references (Ref): (1) - Chiu et al. (2006), (2) - Burgasser et al. (2006b).

survey of UKIDSS and SDSS are listed. In part, they were determined using additional epoch data from 2MASS and DENIS.

For four of the known T dwarfs, these are the first proper motion measurements. For 2MASS J1231+08 and 2MASS J2339+13, we have obtained similar but five to ten times more accurate proper motions than determined earlier by Burgasser et al. (2004) and Burgasser et al. (2003) respectively, even if we prefer the expected proper motion errors ε to our formal errors.


5. Conclusions and discussion

Whereas for three of our candidates we measure moderate $z-J=+2.6...+2.8$ meeting our selection criteria, eight objects have much larger values of $z-J=+3.3...+3.8$ (ULAS J0819+21 has $z-J\approx+3.8$, ULAS J1417+13 and ULAS J2342+08 have $z-J\approx+3.6$) approaching the typical values of T8-T9 dwarfs (Warren et al. 2007; Burningham et al. 2008, 2009; Delorme et al. 2008a).

The mean near-infrared colours $J-K$ and $J-H$ of known T dwarfs follow a clear trend to bluer colour with later subtype (Fig. 3), except for T8-T9 dwarfs which again show colours similar to T6-T7.5 dwarfs. All our new T dwarf candidates exhibit sufficiently blue colours classifying them as $\geq T5$ dwarfs. Based on these colours we assign a preliminary classification as $\approx T5.5$ (with uncertainties up to ± 1.5 subtypes) for ULAS J0819+21, ULAS J0945+07, ULAS J1300+12 (Hip 63510C), and ULAS J1417+13 (both $J-K$ and $J-H$ are between -0.2 and -0.35). For the remaining objects with even larger negative colour indices, including ULAS J1504+05 (= Hip 73786B) with the most extreme values $J-K\approx-0.8$ and $J-H\approx-0.5$, the colours indicate T5-T9 spectral types (Table 7). The larger uncertainty comes here mainly from the already mentioned colour reversal of T8-T9 dwarfs.

Table 7. Estimated spectral types of T dwarf candidates based on colours and absolute magnitudes

Object	SpT ($J-H$)	SpT ($J-K$)	SpT (M_J)	SpT (M_K)	SpT adopted
ULAS J0329+04	T5.0-T9.0 (-0.38)	T5.5-T9.0 (-0.67)			T7.0±2.0
ULAS J0819+21	T5.0-T6.5 (-0.33)	T4.0-T6.5 (-0.23)			T5.5±1.0
ULAS J0945+07	T4.5-T6.0 (-0.22)	T4.0-T7.5 (-0.30)			T5.5±1.5
ULAS J1012+10	T5.0-T9.0 (-0.37)	T5.5-T9.0 (-0.57)			T7.0±2.0
Hip 63510C	T5.0-T6.5 (-0.33)	T4.0-T6.5 (-0.22)	T7.5-T8.0 (16.35)	T6.5-T8.0 (16.56)	T7.0±1.0
ULAS J1417+13	T5.0-T5.5 (-0.23)	T4.0-T6.5 (-0.23)			T5.5±1.0
ULAS J1449+11	T5.0-T9.0 (-0.37)	T5.5-T9.0 (-0.75)			T7.0±2.0
Hip 73786B	T6.5-T9.0 (-0.46)	T6.0-T9.0 (-0.82)	T6.0-T7.5 (15.25)	T6.5-T7.5 (16.07)	T6.5±1.0
ULAS J2320+14	T5.0-T9.0 (-0.34)	T5.0-T9.0 (-0.49)			T7.0±2.0
ULAS J2321+13	T5.5-T9.0 (-0.40)	T5.5-T9.0 (-0.67)			T7.5±1.5
ULAS J2342+08	T5.0-T9.0 (-0.36)	T5.5-T9.0 (-0.61)			T7.0±2.0

**Fig. 3.** Upper panel: Mean $J-K$ vs. $J-H$ (filled symbols) in MKO system for different sub-classes of T dwarfs (labelled; their error bars represent standard deviations) from Leggett et al. (2010) with the colours of our 11 candidates overplotted as crosses with error bars. Lower panel: Zoom to late T dwarf region, where the targets are also labelled.

Absolute magnitudes of the two Hipparcos star companions can be derived using the accurate distances of their primaries. Taking into account the errors in the distances and apparent magnitudes, we get M_J of 16.35 ± 0.04 and 15.25 ± 0.12 , and M_K of 16.56 ± 0.07 and 16.07 ± 0.15 respectively for Hip 63510C and Hip 73786B. These absolute magnitudes place Hip 63510C among the T6-T8 dwarfs with trigonometric parallaxes as given in Leggett et al. (2010), whereas Hip 73786B is as faint as T5.5-T7.5 dwarfs listed in that paper. The fact that the colours hint at a later spectral type for Hip 73786B compared to Hip 63510C, whereas the classification by absolute magnitudes preferred by

us lead to a different conclusion, underlines the need for spectroscopic observations of our new targets and direct comparison with template spectra of the coolest known brown dwarfs. Goldman et al. (2010) discuss Hip 63510C in more detail and come to the conclusion that it is a T8-T9 dwarf with an absolute magnitude and $J-K$ colour pointing to a young age or possible binarity.

Acknowledgements. Data from the UKIDSS 6th data release, SDSS DR7, 2MASS, and DENIS served as the basis for this work. We have also used the SIMBAD and VizieR services at the CDS in Strasbourg and the ARICNS data base on nearby stars in Heidelberg. We thank Doug Finkbeiner for his help accessing the SDSS data at Princeton University and Axel Schwope for comments on a first version of the manuscript. We also thank the referee, Dr. P. Delorme, for helpful comments which led to an improved and extended paper.

References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 54
- Beuzit, J.-L., Segransan, D., Forveille, T., et al. 2004, *A&A*, 425, 997
- Browning, M. K., Basri, G., Marcy, G. W., West, A. A., & Zhang, J. 2010, *AJ*, 139, 504
- Burgasser, A. J., Kirkpatrick, J. D., Brown, M. E., et al. 2002, *ApJ*, 564, 421
- Burgasser, A. J., Kirkpatrick, J. D., McElwain, M. W., Cutri, R. M., Burgasser, A. J., & Skrutskie, M. F. 2003, *AJ*, 125, 850
- Burgasser, A. J., McElwain, M. W., Kirkpatrick, J. D., Cruz, K. L., Tinney, C. G., & Reid, I. N. 2004, *AJ*, 127, 2856
- Burgasser, A. J., Burrows, A., & Kirkpatrick, J. D. 2006a, *ApJ*, 639, 1095
- Burgasser, A. J., Geballe, T. R., Leggett, S. K., Kirkpatrick, J. D., & Golimowski, D. A. 2006b, *ApJ*, 637, 1067
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2008, *MNRAS*, 391, 320
- Burningham, B., Pinfield, D. J., Leggett, S. K., et al. 2009, *MNRAS*, 395, 1237
- Casali, M., Adamson, A., Alves de Oliveira, C., et al. 2007, *A&A*, 467, 777
- Cenarro, A. J., Peletier, R. F., Sánchez-Blázquez, P., et al. 2007, *MNRAS*, 374, 664
- Chiu, K., Fan, X., Leggett, S. K., et al. 2006, *AJ*, 131, 2722
- Delorme, P., Delfosse, X., Albert, L., et al. 2008a, *A&A*, 482, 961
- Delorme, P., Willott, C. J., Forveille, T., et al. 2008b, *A&A*, 484, 469
- Dye, S., Warren, S. J., Hambly, N. C., et al. 2006, *MNRAS*, 372, 1227
- Epchtein, N., de Batz, B., Capoani, L., et al. 1997, *The Messenger*, 87, 27
- Finkbeiner, D. P., Padmanabhan, N., Schlegel, D. J., et al. 2004, *AJ*, 128, 2577
- Gelino, C. R., Kirkpatrick, J. D., & Burgasser, A. J. 2009, online database for 752 L and T dwarfs at DwarfArchives.org (status: 16 November, 2009)
- Goldman, B., Marsat, S., Henning, T., Clemens, C., & Greiner, J. 2010, accepted by *MNRAS*, arXiv:1002.2637
- Hambly, N. C., MacGillivray, H. T., Read, M. A., et al. 2001, *MNRAS*, 326, 1279
- Hambly, N. C., Collins, R. S., Cross, N. J. G., et al. 2008, *MNRAS*, 384, 637
- Heintz, W. D. 1994, *AJ*, 108, 2338
- Hewett, P. C., Warren, S. J., Leggett, S. K., & Hodgkin, S. T. 2006, *MNRAS*, 367, 454
- Irwin et al. 2010, in preparation
- Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, 379, 1599
- Leggett, S. K., Burningham, B., Saumon, D., et al. 2010, *ApJ*, 710, 1627
- Lodieu, N., Pinfield, D. J., Leggett, S. K., et al. 2007, *MNRAS*, 379, 1423
- Lodieu, N., Zapatero Osorio, M. R., Martín, E. L., Solano, E., & Aberasturi, M. 2010, *ApJ*, 708, L107

- Looper, D. L., Kirkpatrick, J. D., & Burgasser, A. J. 2007, *AJ*, 134, 1162
- Pier, J. R.; Munn, Jeffrey A., Hindsley, R. B., et al. 2003, *AJ*, 125, 1559
- Pinfield, D. J., Burningham, B., Tamura, M, et al. 2008, *MNRAS*, 390, 304
- Rauscher, E., & Marcy, G. W. 2006, *PASP*, 118, 617
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Tinney, C. G., Burgasser, A. J., Kirkpatrick, J. D., & McElwain, M. W. 2005, *AJ*, 130, 2326
- Tokunaga, A. T., Simons, D. A., & Vacca, W. D. 2002, *PASP*, 114, 180
- van Leeuwen, F. 2007, *A&A*, 474, 653
- Warren, S. J., Mortlock, D. J., Leggett, S. K., et al. 2007, *MNRAS*, 381, 1400